

INSTRUCTIONS TO PATIENTS FOLLOWING ORAL SURGERY

1. Bite on gauze for 1/2 hour
2. Keep fingers and tongue away from operated area.
3. Do not rinse your mouth for 24 hours, although it is permissible to drink cool and lukewarm liquids. Smoking should be avoided for 12 hours
4. Starting tomorrow, rinse your mouth frequently with a solution of ½ teaspoon of salt in a glass of warm water. Continue the rinses for a few days.
5. If you have any discomfort, take whatever you normally take for a headache, every few hours. If necessary, Dr. Xiang will prescribe stronger medication for you.
6. Diet – cold or lukewarm liquids may be taken for the first 4-6 hours. After this any soft food is permissible.
7. Bleeding – it is normal for the saliva to be streaked with blood for a day. If frank bleeding is present, fold damp, sterile gauze into a firm wad and place it directly on the bleeding area. Maintain firm pressure by biting for 20 minutes. The gauze may be substituted by a teabag. The tannic acid in the tea has a clotting effect.
8. Swelling and discoloration is to be expected in certain areas, usually reaching its maximum two days after surgery. It will disappear gradually and is no cause for concern. If desired, ice packs may be applied for the first 4-6 hours only, alternating 10 minutes on, 10 minutes off.
9. Do not hesitate to call the office if in doubt.

Dr. Peter X. Xiang DMD, PhD, M.Dent, FRDC (C)

Telephone: 204-726-1211 Office
204-891-5767 Cell